


PROFILE FOR THORNEY MEDICAL CENTRE COMPARED WITH Peterborough PCT

APHO General Practice Profiles



Practice Peterborough PCT Peterborough PCT
 Average Total

Population 7049 6586 177832

Source: QOF 2009/10

Demography and patient satisfaction

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|----------------|-----------|-----------------|----------------|---------------|-----------------|
| % aged 0 to 4 years | 5.3% | 7.5% | 5.9% | 0.0% | | 16.9% |
| % aged 5 to 14 years | 11.1% | 12.5% | 11.2% | 0.0% | | 28.9% |
| % aged under 15 years | 16.4% | 19.9% | 17.1% | 0.0% | | 42.6% |
| % aged 65+ years | 18.1% | 12.6% | 15.8% | 0.0% | | 45.2% |
| % aged 75+ years | 8.2% | 6.2% | 7.5% | 0.0% | | 26.5% |
| % aged 85+ years | 2.3% | 1.7% | 2.1% | 0.0% | | 12.0% |
| Deprivation score (IMD 2007) | 18.5 | 24.5 | 21.7 | 2.6 | | 68.9 |
| IDACI (Income Deprivation Affecting Children) | 0.15 | - | 0.24 | 0.03 | | 0.80 |
| IDAOPPI (Income Deprivation Affecting Older People) | 0.17 | - | 0.22 | 0.05 | | 0.74 |
| % satisfied with phone access | 74.3% | 58.8% | 64.7% | 12.1% | | 96.2% |
| % able to see a doctor within 2 days | 91.7% | 81.4% | 80.1% | 0.0% | | 100.0% |
| % able to book appointment >= 2d ahead | 51.3% | 65.7% | 69.5% | 0.0% | | 100.0% |
| % satisfied with opening hours | 81.5% | 81.1% | 79.4% | 43.5% | | 100.0% |
| % able to see preferred GP | 80.8% | 69.1% | 71.4% | 0.0% | | 100.0% |

Cardiovascular disease - Coronary heart disease

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|----------------|-----------|-----------------|----------------|---------------|-----------------|
| CHD: QOF prevalence (all ages) | 3.7% | 2.9% | 3.4% | 0.0% | | 9.8% |
| Heart failure w LVD: QOF prevalence | 0.3% | 0.3% | 0.4% | 0.0% | | 2.4% |
| Exception rate for CHD indicators | 5.9% | 9.3% | 7.7% | 0.0% | | 50.0% |
| CHD Emergency Admission Rate | 1.9 | 2.1 | 2.5 | 0.0 | | 13.9 |
| CHD Elective Admission Rate | 3.9 | 2.3 | 1.8 | 0.0 | | 10.0 |
| Ratio of recorded vs expected CHD prevalence | 0.8 | 0.8 | 0.8 | 0.0 | | 2.8 |
| CHD 2: Angina referred for exercise testing &/or ass... | 93.8% | 94.3% | 94.7% | 0.0% | | 100.0% |
| CHD 5: Record of BP in the previous 15mths | 98.5% | 97.5% | 97.7% | 64.0% | | 100.0% |
| CHD 6: Last BP reading in last 15mths is <=150/90 | 93.1% | 89.0% | 89.8% | 0.0% | | 100.0% |
| CHD 7: Record of total cholesterol in last 15mths | 92.3% | 92.4% | 93.7% | 45.8% | | 100.0% |
| CHD 8: Last total cholesterol is <=5mmol/l | 84.8% | 81.3% | 82.1% | 31.0% | | 100.0% |
| CHD 9: Record that aspirin, APT or ACT is taken | 94.9% | 92.8% | 93.9% | 0.0% | | 100.0% |
| CHD 10: Currently treated with beta blocker | 68.4% | 73.1% | 73.7% | 28.6% | | 100.0% |
| CHD 11: History of MI: treated with ACE-I | 86.8% | 87.3% | 89.1% | 0.0% | | 100.0% |
| CHD 12: Influenza immunisation given 1 Sep - 31 Mar | 92.1% | 91.9% | 91.9% | 0.0% | | 100.0% |

Cardiovascular disease - Stroke and TIA

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Stroke: QOF prevalence (all ages) | 1.9% | 1.4% | 1.7% | 0.0% | | 6.0% |
| Exception rate for stroke indicators | 7.4% | 8.0% | 7.1% | 0.0% | | 100.0% |
| Stroke 13: New patients referred for further investi... | 100.0% | 89.1% | 90.2% | 0.0% | | 100.0% |
| Stroke 5: BP recorded in last 15mths | 98.4% | 96.4% | 96.8% | 0.0% | | 100.0% |
| Stroke 6: Last BP reading is 150/90 or less | 92.2% | 87.3% | 88.1% | 50.0% | | 100.0% |
| Stroke 7: Total cholesterol recorded in last 15mths | 92.7% | 90.5% | 91.4% | 0.0% | | 100.0% |
| Stroke 8: Last measured total cholesterol <=5mmol/l | 78.0% | 78.1% | 77.3% | 0.0% | | 100.0% |
| Stroke 12: Record of aspirin, APT or ACT taken | 97.2% | 94.6% | 94.1% | 0.0% | | 100.0% |
| Stroke 10: Influenza immunisation given 1 Sep-31 Mar | 90.5% | 89.3% | 89.0% | 0.0% | | 100.0% |

Cardiovascular disease - Heart failure and atrial fibrillation

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Heart failure: QOF prevalence (all ages) | 0.8% | 0.6% | 0.7% | 0.0% | | 3.8% |
| Heart failure w LVD: QOF prevalence | 0.3% | 0.3% | 0.4% | 0.0% | | 2.4% |
| Exception rate for heart failure indicators | 25.9% | 21.2% | 17.2% | 0.0% | | 100.0% |
| Atrial fibrillation: QOF prevalence | 1.1% | 1.0% | 1.4% | 0.0% | | 5.0% |
| Exception rate for atrial fibrillation indicators | 4.3% | 4.0% | 4.1% | 0.0% | | 66.7% |
| HF 2: Diagnosis conf. by ECG/specialist assessm. | 90.9% | 96.6% | 95.9% | 0.0% | | 100.0% |
| HF 3: HF w LVD: treated with ACE-I or ARB | 85.0% | 89.7% | 95.9% | 0.0% | | 100.0% |
| HF 4: HF w LVD: ACE inh. or ARB and beta-blocker | 66.7% | 81.0% | 82.4% | 0.0% | | 100.0% |
| AF 4: diagnosed with ECG or by specialist | 100.0% | 96.6% | 96.5% | 0.0% | | 100.0% |
| AF 3: treated w anti-coag./platelet therapy | 91.5% | 92.6% | 93.6% | 0.0% | | 100.0% |

Cardiovascular disease - Risk factors for CVD

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|--|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Hypertension: QOF prevalence (all ages) | 15.6% | 12.4% | 13.4% | 0.1% | | 36.7% |
| Exception rate for hypertension indicators | 3.7% | 4.6% | 4.9% | 0.0% | | 51.9% |
| Exception rate for smoking indicators | 0.4% | 0.6% | 0.7% | 0.0% | | 13.7% |
| Obesity: QOF prevalence (16+) | 7.6% | 11.3% | 10.5% | 0.5% | | 29.7% |
| BP 4: Record of BP in last 9mths | 92.7% | 91.5% | 91.5% | 45.3% | | 100.0% |
| BP 5: Last (9mths) blood pressure <=150/90 | 84.2% | 78.0% | 78.7% | 39.7% | | 100.0% |
| PP 1: CV risk assessment for new hypertension cases | 91.2% | 81.3% | 81.7% | 0.0% | | 100.0% |
| PP 2: life style advice for new hypertension cases | 92.1% | 85.4% | 84.5% | 0.0% | | 100.0% |
| Dep 1: Depression case finding in CHD&/diabetes pats | 93.9% | 89.0% | 88.3% | 0.0% | | 100.0% |
| Smoking 3: status recorded in last 15mths | 97.3% | 96.0% | 95.2% | 58.4% | | 100.0% |
| Smoking 4: cessation advice/referral offered | 92.1% | 92.9% | 92.8% | 6.7% | | 100.0% |

Diabetes

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Diabetes: QOF prevalence (17+) | 5.8% | 5.6% | 5.4% | 0.2% | | 14.3% |
| Exception rate for diabetes indicators | 6.5% | 8.6% | 6.4% | 0.1% | | 36.6% |
| Hypertension: QOF prevalence (all ages) | 15.6% | 12.4% | 13.4% | 0.1% | | 36.7% |
| Exception rate for hypertension indicators | 3.7% | 4.6% | 4.9% | 0.0% | | 51.9% |
| Exception rate for smoking indicators | 0.4% | 0.6% | 0.7% | 0.0% | | 13.7% |
| Obesity: QOF prevalence (16+) | 7.6% | 11.3% | 10.5% | 0.5% | | 29.7% |
| Diabetes admission rate | 0.9 | 1.1 | 1.1 | 0.0 | | 10.9 |
| Ratio of recorded vs expected diabetes prevalence | 0.8 | 1.0 | 0.9 | 0.2 | | 5.8 |
| DM 2: Record of BMI in the last 15mths | 96.0% | 94.8% | 94.7% | 52.9% | | 100.0% |
| DM 5: Record of HbA1c/equivalent last 15mths | 98.4% | 97.5% | 97.2% | 57.9% | | 100.0% |
| DM 23: Last HbA1c is <=7 in last 15mths | 45.6% | 47.9% | 53.8% | 0.0% | | 100.0% |
| DM 24: Last HbA1c is <=8 in last 15mths | 79.1% | 74.9% | 77.4% | 9.3% | | 100.0% |
| DM 25: Last HbA1c is <=9 in last 15mths | 91.2% | 86.9% | 88.0% | 9.3% | | 100.0% |
| DM 21: Retinal screening in last 15mths | 91.9% | 89.6% | 90.9% | 3.4% | | 100.0% |
| DM 9: Record of peripheral pulses last 15mths | 95.3% | 92.4% | 91.1% | 11.3% | | 100.0% |
| DM 10: Record of neuropathy test last 15mths | 94.9% | 92.1% | 90.8% | 17.0% | | 100.0% |
| DM 11: Record of BP in last 15mths | 98.5% | 98.6% | 98.3% | 78.0% | | 100.0% |
| DM 12: Last BP is <=145/85 | 82.8% | 79.1% | 80.6% | 41.9% | | 100.0% |
| BP 4: Record of BP in last 9mths | 92.7% | 91.5% | 91.5% | 45.3% | | 100.0% |
| BP 5: Last (9mths) blood pressure <=150/90 | 84.2% | 78.0% | 78.7% | 39.7% | | 100.0% |
| DM 13: Record of micro-albuminuria test last 15mths | 91.3% | 88.8% | 88.5% | 1.0% | | 100.0% |
| DM 22: eGRF or serum creatinin testing in last 15mths | 99.4% | 97.6% | 97.0% | 55.1% | | 100.0% |
| DM 15: Proteinuria/micro-album. treated w inhibitors | 83.3% | 83.9% | 88.8% | 0.0% | | 100.0% |
| DM 16: Record of total cholesterol last 15mths | 98.1% | 96.7% | 96.0% | 60.8% | | 100.0% |
| DM 17: Measured total chol (last 15mths) <=5mmol/l | 87.6% | 83.6% | 83.0% | 46.2% | | 100.0% |
| DM 18: Influenza immunisation given 1 Sep - 31 Mar | 93.5% | 91.2% | 90.4% | 15.1% | | 100.0% |
| Dep 1: Depression case finding in CHD&/diabetes pats | 93.9% | 89.0% | 88.3% | 0.0% | | 100.0% |
| Smoking 3: status recorded in last 15mths | 97.3% | 96.0% | 95.2% | 58.4% | | 100.0% |
| Smoking 4: cessation advice/referral offered | 92.1% | 92.9% | 92.8% | 6.7% | | 100.0% |

Mental health

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Psychoses: QOF prevalence (all ages) | 0.5% | 0.8% | 0.8% | 0.0% | | 30.5% |
| Exception rate for patients with psychoses indicators | 8.5% | 8.0% | 10.8% | 0.0% | | 61.5% |
| Dementia: QOF prevalence (all ages) | 0.7% | 0.3% | 0.5% | 0.0% | | 8.7% |
| Exception rate for dementia indicators | 2.1% | 5.0% | 7.1% | 0.0% | | 100.0% |
| Depression: QOF prevalence (18+) | 17.5% | 9.9% | 10.9% | 0.0% | | 47.5% |
| Exception rate for depression indicators | 3.7% | 5.7% | 6.0% | 0.0% | | 64.3% |
| Learning disability: QOF prevalence (18+) | 0.5% | 0.3% | 0.4% | 0.0% | | 6.0% |
| MH 9: full review in last 15mths | 92.3% | 91.2% | 92.2% | 0.0% | | 100.0% |
| MH 4: Li-therapy: record of s.creat. & TSH last 15mths | 100.0% | 99.2% | 97.6% | 0.0% | | 100.0% |
| MH 5: Lithium therapy: record of Li-levels last 6mths | 100.0% | 86.2% | 91.2% | 0.0% | | 100.0% |
| MH 6: with comprehensive care plan | 88.9% | 89.3% | 88.9% | 0.0% | | 100.0% |
| MH 7: Psychoses pat. who dna AR: followed up within 14d | 100.0% | 92.7% | 93.9% | 0.0% | | 100.0% |
| Dem 2: Dementia care has been reviewed last 15mths | 87.0% | 80.0% | 79.3% | 0.0% | | 100.0% |
| Dep 1: Depression case finding in CHD&/diabetes pats | 93.9% | 89.0% | 88.3% | 0.0% | | 100.0% |
| Dep 2: Depression cases with severity assessment | 97.7% | 90.2% | 91.8% | 0.0% | | 100.0% |
| Dep 3: 2 severity assessm. for new depression cases | 92.9% | 67.5% | 70.1% | 0.0% | | 100.0% |

Respiratory disease

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|----------------|-----------|-----------------|----------------|---------------|-----------------|
| COPD: QOF prevalence (all ages) | 1.1% | 1.4% | 1.6% | 0.0% | | 7.3% |
| Exception rate for COPD indicators | 9.6% | 12.7% | 12.6% | 0.0% | | 100.0% |
| COPD admission rate | 2.2 | 1.8 | 2.0 | 0.0 | | 17.2 |
| Asthma: QOF prevalence (all ages) | 6.6% | 5.7% | 5.9% | 1.1% | | 13.2% |
| Exception rate for asthma indicators | 2.0% | 4.7% | 5.2% | 0.0% | | 51.7% |
| Exception rate for smoking indicators | 0.4% | 0.6% | 0.7% | 0.0% | | 13.7% |
| COPD 12: Diagnosis conf. by spirometry in last 15mths | 100.0% | 87.9% | 90.3% | 0.0% | | 100.0% |
| COPD 10: Record of FeV1 in last 15mths | 89.0% | 90.1% | 88.5% | 0.0% | | 100.0% |
| COPD 13: assessed using MRC dyspnoea score last 15mths | 90.1% | 87.0% | 89.9% | 0.0% | | 100.0% |
| COPD 8: Influenza immunisation given 1 Sep - 31 Mar | 93.2% | 93.0% | 92.7% | 0.0% | | 100.0% |
| Asthma 8: with measures of variability/reversibility... | 100.0% | 86.9% | 87.6% | 0.0% | | 100.0% |
| Asthma 3: smoking recorded in last 15 mths (14-19y) | 89.2% | 89.6% | 89.0% | 33.3% | | 100.0% |
| Asthma 6: review in the last 15mths | 75.0% | 79.7% | 78.0% | 0.0% | | 100.0% |
| Smoking 3: status recorded in last 15mths | 97.3% | 96.0% | 95.2% | 58.4% | | 100.0% |
| Smoking 4: cessation advice/referral offered | 92.1% | 92.9% | 92.8% | 6.7% | | 100.0% |

Secondary Care Use

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|----------------|-----------|-----------------|----------------|---------------|-----------------|
| All GP referrals (1st attendance) | 179.5 | 218.6 | 186.8 | 34.0 | | 615.0 |
| All attendances | 792.9 | 805.1 | 739.9 | 105.2 | | 1581.5 |
| Dermatology, GP referrals (1st attendance) | 8.6 | 10.1 | 13.7 | 0.0 | | 71.8 |
| General Medicine, GP referrals (1st attendance) | 5.9 | 5.9 | 10.9 | 0.0 | | 81.3 |
| General Medicine - return ratio | 1.7 | 1.8 | 2.7 | 0.5 | | 31.0 |
| General Surgery, GP referrals (1st attendance) | 16.5 | 22.5 | 23.1 | 1.9 | | 70.8 |
| General Surgery - return ratio | 1.6 | 1.3 | 1.5 | 0.3 | | 5.1 |
| Gynaecology, GP referrals (1st attendance) | 17.8 | 24.3 | 17.1 | 2.2 | | 56.4 |
| Gynaecology - return ratio | 1.6 | 1.6 | 1.3 | 0.2 | | 8.5 |
| Orthopaedics, GP referrals (1st attendance) | 21.9 | 19.8 | 20.2 | 0.0 | | 81.4 |
| Orthopaedics - return ratio | 2.2 | 2.1 | 1.8 | 0.7 | | 5.0 |
| Paediatrics, GP referrals (1st attendance) | 5.3 | 10.7 | 5.7 | 0.0 | | 34.6 |
| Urology, GP Referrals (first attendance) | 6.3 | 8.5 | 7.9 | 0.0 | | 29.7 |
| Urology - return ratio | 2.1 | 1.7 | 2.1 | 0.3 | | 20.5 |
| A&E attendances per 1000 population | 169.5 | 224.2 | 372.8 | 8.6 | | 1190.8 |
| All admissions - All ages | 217.5 | 232.8 | 231.6 | 26.5 | | 503.0 |
| All admissions - A&E | 9.2 | 10.0 | 9.6 | 0.0 | | 84.5 |
| All admissions - CHD | 5.9 | 4.8 | 4.7 | 0.0 | | 25.9 |
| All admissions - Respiratory | 16.1 | 17.8 | 16.0 | 1.2 | | 51.7 |
| Elective Admission Rate | 114.3 | 106.7 | 120.6 | 10.5 | | 354.4 |
| Emergency Admission Rate | 81.5 | 89.6 | 86.7 | 10.7 | | 267.4 |
| Ambulatory Care Sensitive Conditions Rate | 16.0 | 16.5 | 14.5 | 1.5 | | 61.0 |
| CHD Elective Admission Rate | 3.9 | 2.3 | 1.8 | 0.0 | | 10.0 |
| CHD Emergency Admission Rate | 1.9 | 2.1 | 2.5 | 0.0 | | 13.9 |
| Diabetes admission rate | 0.9 | 1.1 | 1.1 | 0.0 | | 10.9 |
| COPD admission rate | 2.2 | 1.8 | 2.0 | 0.0 | | 17.2 |
| Cancer admission rate | 33.9 | 22.3 | 28.1 | 1.3 | | 105.2 |
| Emergency readmissions within 28d of discharge | 5.6% | 7.5% | 7.7% | 0.0% | | 31.3% |

Other conditions

| Indicator | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Cancer: QOF prevalence (all ages) | 1.8% | 1.2% | 1.4% | 0.0% | | 3.9% |
| Exception rate for cancer indicators | 4.2% | 1.2% | 1.5% | 0.0% | | 100.0% |
| Cancer admission rate | 33.9 | 22.3 | 28.1 | 1.3 | | 105.2 |
| CKD: QOF prevalence (18+) | 4.1% | 3.8% | 4.3% | 0.0% | | 19.8% |
| Exception rate for CKD indicators | 2.5% | 2.2% | 4.3% | 0.0% | | 100.0% |
| Epilepsy: QOF prevalence (18+) | 0.9% | 0.8% | 0.8% | 0.1% | | 3.8% |
| Exception rate for epilepsy indicators | 10.0% | 8.6% | 7.8% | 0.0% | | 66.7% |
| Hypothyroidism: QOF prevalence (all ages) | 3.8% | 2.9% | 2.9% | 0.0% | | 7.9% |
| Exception rate for hypothyroidism indicators | 1.5% | 0.5% | 0.5% | 0.0% | | 17.9% |
| Palliative/supportive care: QOF prevalence (all ages) | 0.1% | 0.1% | 0.1% | 0.0% | | 2.3% |
| Cancer 3: review within 6mths of diagnosis | 91.3% | 94.3% | 93.1% | 0.0% | | 100.0% |
| CKD 2: Record of BP in last 15mths | 97.3% | 97.7% | 97.6% | 64.7% | | 100.0% |
| CKD 3: Last BP reading measured in last 15mths is <=... | 71.0% | 71.1% | 73.9% | 29.4% | | 100.0% |
| CKD 5: Hypertension treated with ACE inhibitor/ARB | 100.0% | 92.0% | 91.8% | 0.0% | | 100.0% |
| CKD 6: Urine albumin : creatinine ratio test last 15... | 83.0% | 80.3% | 77.7% | 0.0% | | 100.0% |
| Epilepsy 6: Record of seizure frequency | 95.8% | 96.1% | 95.3% | 0.0% | | 100.0% |
| Epilepsy 7: Medication review w patient/carer last 1... | 95.8% | 95.6% | 95.0% | 0.0% | | 100.0% |
| Epilepsy 8: Seizure free for last 12mths | 82.1% | 75.3% | 74.4% | 0.0% | | 100.0% |
| Thyroid 2: function test recorded last 15mths | 93.9% | 95.5% | 95.8% | 50.0% | | 100.0% |

Population

This section of the Profiles provides an overview of the practice(s) and their populations, by age and sex, and by deprivation. This information is essential in understanding the very basic needs of the local population. Data on access and patient satisfaction come from the GP Patient Survey.

Data sources for GP practice populations vary substantially and there is little consensus on which is the most reliable source. The figure displayed here is from QOF. The criteria for exclusion of a practice and a detailed explanation of how we dealt with these issues can be found in the metadata document.

Indicators

The light grey bar shows the range of values found in England. The dark grey sections mark out the range within which the middle-half of the observed values lie (25th to 75th percentile). The red line shows where the England average is. The position of the dot shows the practice value, a diamond the PCT value, a triangle the cluster value in relation to this scale. The corresponding numbers can be found in the cells next to the chart.

The practice values are colour-coded:

- Significantly different from national average
- Not significantly different from regional average
- Significance level not calculated
- PCT value
- Cluster value

